195 mm

275 mm

PATIENT INFORMATION LEAFLET

Scheduling status:
S0

SEPTOGARD® lozenges
Mentha piperata (peppermint, crystals); *Glycyrrhiza glabra* L. (liquorice, root); *Terminalia belerica* Roxb. (Beleric myrobalan, fruit); *Zingiber officinale* Rosc. (ginger, rhizome); *Curcuma longa* L. (turmeric, rhizome).
Contains sugar: sucrose 1375 mg/lozenge, liquid glucose 1375 mg/lozenge.
Contains preservatives: potassium sorbate 0,2 % m/m, sorbic acid 0,02 % m/m.
• D33.8. Complementary Medicine: Discipline-specific. Other – herbal.
• This unregistered medicine has not been evaluated by the SAHPRA for its quality, safety or intended use.

Read all of this leaflet carefully because it contains important information for you
SEPTOGARD® lozenges is available without a doctor's prescription. Nevertheless, you still need to use SEPTOGARD® lozenges carefully to get the best results from it.
• Keep this leaflet. You may need to read it again.
• Do not share SEPTOGARD® lozenges with any other person.
• Ask your healthcare provider or pharmacist if you need more information or advice.
• You must see a doctor if your symptoms worsen or do not improve after 5 days.

What is in this leaflet

1. What SEPTOGARD® lozenges is and what it is used for
2. What you need to know before you use SEPTOGARD® lozenges
3. How to use SEPTOGARD® lozenges
4. Possible side effects
5. How to store SEPTOGARD® lozenges
6. Contents of the pack and other information

1. What SEPTOGARD® lozenges is and what it is used for
SEPTOGARD® lozenges is a herbal medicine that contains ingredients which can assist with the symptomatic relief of sore throat, cough and congestion. Efficacy of support may vary between users.

2. What you need to know before you use SEPTOGARD® lozenges
Do not use SEPTOGARD® lozenges:
• If you are hypersensitive (allergic) towards any of the ingredients of SEPTOGARD® lozenges (listed in section 6 **Contents of pack and other information**).
• If you have epilepsy, asthma, persistent or chronic cough or other chronic lung conditions, as SEPTOGARD® lozenges contains peppermint.
• If you have high blood pressure or are taking medicine for high blood pressure (see **Other medicines and SEPTOGARD® lozenges**).
• If you have low blood potassium levels or are taking diuretics (water tablets), cardiac glycosides, corticosteroids, laxatives or medications which may aggravate electrolyte imbalance (see **Other medicines and SEPTOGARD® lozenges**).
• Do not consume with other liquorice-containing products.
• If you have a heart, kidney or liver disorder.
• If you are pregnant or breastfeeding (see **Pregnancy, breastfeeding and fertility**).
• SEPTOGARD® lozenges contains ingredients that are not suitable for use in children under the age of 4 years.
• If you are taking any chronic medication, do not use this product without consulting your medical practitioner.

Warnings and precautions
Special care should be taken with SEPTOGARD® lozenges.
• Consult a healthcare provider or doctor before use if you have/are taking:
• a bleeding disorder or blood thinners (anticoagulant/antiplatelet medication) [see **Other medicines and SEPTOGARD® lozenges**]; or
• (a history of) gallstones, bile duct obstruction, stomach ulcers or excess stomach acid as SEPTOGARD® lozenges contains turmeric rhizome.
• If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before using SEPTOGARD® lozenges.
• SEPTOGARD® lozenges contains glucose and sucrose which may have an effect on the control of your blood sugar if you have diabetes mellitus.
• Discontinue use and consult a healthcare provider or doctor if your sore throat symptoms worsen or continue for more than 2 days.
• Discontinue use and consult a healthcare provider or doctor if your cough worsens and continues for more than 7 days or is accompanied by a fever.
• Discontinue use at least 2 weeks before any surgical procedures.
• Do not exceed the daily recommended dose.
• Do not use after the date of expiry.
• Products containing herbal ingredients may interact with some medication.
Porphyria: Safety has not been established.
Tell your doctor if you are not sure about any of the above.

Other medicines and SEPTOGARD® lozenges
Always tell your healthcare provider if you are taking any other medicine, including complementary or traditional medicines. Consult your doctor, pharmacist or healthcare provider before use if you are taking:
• medication for high blood pressure, as SEPTOGARD® lozenges contains ingredients which may influence the efficacy of these medicines;
• blood thinners (anticoagulant or antiplatelet medication) e.g., warfarin, as SEPTOGARD® lozenges contains some ingredients which may influence the efficacy of these medicines;
• cardiac glycosides (heart medication) e.g., digoxin, as liquorice root may increase the risk of cardiotoxicity;
• corticosteroids, as liquorice root can increase the effects and side effects of corticosteroids;
• sulfasalazine (for arthritis), as turmeric rhizome may increase the effects and side effects of sulfasalazine; and
• thiazide diuretics (water tablets), stimulant laxatives or other medications which may aggravate electrolyte imbalance, as liquorice root can increase potassium loss.

Using SEPTOGARD® lozenges with food, drink and alcohol
There is no known interaction between the contents of SEPTOGARD® lozenges and food, drink or alcohol.

Pregnancy, breastfeeding and fertility
Do not use SEPTOGARD® lozenges while you are pregnant or breastfeeding as it contains liquorice root and beleric myrobalan fruit which are contraindicated. If you are pregnant or breastfeeding, think you may be pregnant or are planning to have a baby, please consult your doctor, pharmacist, or other healthcare provider for advice before taking this product.

Driving and using machines
SEPTOGARD® lozenges should not affect your ability to drive or use machinery. However, it is not always possible to predict to what extent SEPTOGARD® lozenges may interfere with your daily activities. Ensure that you do not engage in the above activities until you are aware of the measure to which SEPTOGARD® lozenges affects you.

SEPTOGARD® lozenges contains glucose and sucrose
If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before using SEPTOGARD® lozenges.

3. How to use SEPTOGARD® lozenges
Do not share medicines with any other person.
Always use SEPTOGARD® lozenges exactly as described in this leaflet or as your doctor, pharmacist or nurse has told you. You should check with your doctor, pharmacist or nurse if you are unsure.

The usual dose for adults and children 4 years and older is: Dissolve one (1) lozenge slowly in the mouth, every three to four hours. Do not exceed six (6) lozenges per 24 hours. For best results, do not to eat or drink anything within 10 minutes of consuming a lozenge.

If you use more SEPTOGARD® lozenges than you should
In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or Poisons Information Centre.

If you forget to use SEPTOGARD® lozenges
Use your missed dose when you remember. If it is almost time for your next dose, skip the missed dose and consume SEPTOGARD® lozenges at the next regularly scheduled time. Do not consume a double dose to make up for forgotten individual doses.

If you stop using SEPTOGARD® lozenges
If you are using the dose as indicated, it is unlikely that there should be withdrawal effects when you stop using this medication.

4. Possible side effects
SEPTOGARD® lozenges can have side effects. Not all side effects reported for SEPTOGARD® lozenges are included in this leaflet. If your general health worsens or if you experience any untoward effects while using SEPTOGARD® lozenges, please consult your healthcare provider for advice.

If any of the following or other side effects occur, stop taking this product and consult your doctor or pharmacist or go to the casualty department at your nearest hospital:
Allergic reactions such as: skin rashes, hives (skin bumps), itchiness, swelling of the face, lips mouth or throat, which may cause difficulty in swallowing or breathing. These are all very serious side effects. If you have them, you may have had a serious reaction to SEPTOGARD® lozenges. You may need urgent medical attention or hospitalisation.

Tell your doctor or pharmacist if you notice any of the following:
Frequency not known:
• Headache.
• Abdominal pain or discomfort, anal burning, belching, bloating, burning aftertaste, constipation, diarrhoea, dry mouth, dyspepsia, heartburn/reflux, nausea and vomiting.

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

Reporting of side effects
If you get side effects, talk to your doctor, pharmacist or nurse. You can also report side effects to SAHPRA via the Med Safety APP (Medsafety X SAHPRA) and eReporting platform (who-umc.org), found on SAHPRA's website. By reporting side effects, you can help provide more information on the safety of SEPTOGARD® lozenges.

5. How to store SEPTOGARD® lozenges
• Store all medicines out of reach of children.
• Store in a dry place at or below 25 °C.
• Protect from light and moisture.
• Keep in original packaging until use.
• Do not refrigerate.
• Do not store in a bathroom.
• Do not use after the expiry date stated on the label and carton.
• Return all unused medicine to your pharmacist.
• Do not dispose of unused medicine in drains or sewerage systems (e.g., toilets).

6. Contents of pack and other information
What SEPTOGARD® lozenges contains

Each Lozenge contains:	Per lozenge	Per maximum daily dose (6 lozenges)
Mentha piperata (peppermint/Sanskrit – Pudina Ke Phool) [crystals]	7 mg	42 mg
Extract of the following:		
Glycyrrhiza glabra L. (liquorice/Sanskrit - Yashtimadhu) [root]	20 mg	120 mg
Terminalia belerica Roxb. (beleric myrobalan/Sanskrit – Baheda) [fruit]	5,5 mg	33 mg
Zingiber officinale Rosc. (ginger/Sanskrit – Sonth) [rhizome]	4,5 mg	27 mg
Curcuma longa L. (turmeric/Sanskrit – Haldi) [rhizome]	2 mg	12 mg

The other ingredients and excipients are: citric acid monohydrate.
Honey-Lemon flavour: honey-lemon flavouring, quinoline yellow colourant.
Mint flavour: propylene glycol, brilliant blue colourant, quinoline yellow colourant.
Orange flavour: orange flavouring, sunset yellow colourant.

SEPTOGARD® lozenges contains sugar: sucrose 1375 mg/lozenge, liquid glucose 1375 mg/lozenge.
SEPTOGARD® lozenges contains preservatives: potassium sorbate 0,2 % m/m, sorbic acid 0,02 % m/m.

What SEPTOGARD® lozenges looks like and contents of the pack
SEPTOGARD® lozenges are golden-yellow, green or orange coloured, flat, bevelled, circular lozenges with a methanolic sweet taste, characteristic of either honey-lemon, mint, or orange flavour.
SEPTOGARD® lozenges are available as 12 lozenges (containing 2 blister packs of 6 lozenges each) or 24 lozenges (containing 4 blister packs of 6 lozenges each) in a printed unit carton that includes a patient information leaflet.

Holder of Certificate of Registration
Nativa (Pty) Ltd
260 Cradock Avenue, Lyttelton, Centurion, 0157, Gauteng, South Africa
Tel: +27 (0) 12 664 7110
Customer care line: 0860 628 482
Email: health@nativa.co.za
Website: www.nativa.co.za

This leaflet was last revised in
April 2025

Date of registration
To be allocated

Registration number
To be allocated

Professional Information
The SEPTOGARD® lozenges Professional Information may be obtained from the Nativa website (www.nativa.co.za).

PPAT155/02

PEI0465 0425 E

195 mm

275 mm

PASIËNTINLIGTINGSBROSJURE

Skeduleringstatus:
S0

SEPTOGARD® suigtablette
Mentha piperata (peperment, kristalle); *Glycyrrhiza glabra* L. (dropwortel); *Terminalia belerica* Roxb. (beleriese myrobalan-vrug); *Zingiber officinale* Rosc. (gemmer-risoom); *Curcuma longa* L. (borrie-risoom).
Bevat suiker: sukrose 1375 mg/suigtablet, vloeibare glukose 1375 mg/suigtablet.
Bevat preserveermiddels: kaliumsorbaat 0,2 % m/m, sorbiensuur 0,02 % m/m.
• D33.8. Komplementêre Medisyne: Dissipliene-spesifiek. Ander – kruie.
• Hierdie medisyne is nog nie deur die SAHPRA geregistreer of geëvalueer vir kwaliteit, veiligheid of voorgestelde gebruik nie.

Lees die hele brosjure deeglik want dit bevat belangrike inligting
SEPTOGARD® suigtablette is beskikbaar sonder 'n dokter se voorskrif. U moet egter steeds SEPTOGARD® suigtablette met omsigtigheid gebruik om die beste resultate daarmee te verkry.
• Hou hierdie brosjure. Dit kan nodig wees dat u dit weer moet lees.
• Moet nie SEPTOGARD® suigtablette met enige ander persoon deel nie.
• Indien u verdere inligting of advies verlang, vra asseblief u gesondheidsorgverskaffer of apteker.
• Raadpleeg 'n dokter indien u simptome vererger of nie verbeter na 5 dae nie.

Wat is in hierdie brosjure

1. Wat SEPTOGARD® suigtablette is en waarvoor dit gebruik word
2. Wat u moet weet voordat u SEPTOGARD® suigtablette gebruik
3. Hoe om SEPTOGARD® suigtablette te gebruik
4. Moontlike newe-effekte
5. Hoe om SEPTOGARD® suigtablette te bewaar
6. Inhoud van die verpakking en ander inligting

1. Wat SEPTOGARD® suigtablette is en waarvoor dit gebruik word
SEPTOGARD® suigtablette is 'n kruiemedisyne wat bestanddele bevat wat simptomatiese verligting kan bied vir seerkeel, hoes en verstopping. Effektiwiteit van ondersteuning kan tussen gebruikers verskil.

2. Wat u moet weet voordat u SEPTOGARD® lozenges gebruik
Moet nie SEPTOGARD® suigtablette gebruik nie:
• Indien u hipersensitief (allergies) is vir enige van die bestanddele in SEPTOGARD® suigtablette (gelys in afdeling 6 **Inhoud van die verpakking en ander inligting**).
• Indien u epilepsie, asma, aanhoudend of chronies hoes of ander chroniese longtoestande het, aangesien SEPTOGARD® suigtablette peperment bevat.
• Indien u hoë bloeddruk het of medikasie vir hoë bloeddruk neem (sien **Ander medikasie en SEPTOGARD® suigtablette**).
• Indien u lae kaliumbloedvlakke het, of diuretikums (waterpille), hartglikosiede, kortikosteroïede, lakseermiddels of medikasie wat elektrolietbalans kan beïnvloed, neem (sien **Ander medikasie en SEPTOGARD® suigtablette**).
• Moet nie saam met ander drop-bevattende produkte gebruik nie.
• Indien u 'n hart-, nier- of lewerversteuring het.
• Indien u swanger is of borsvoed (sien **Swangerskap, borsvoeding en vrugbaarheid**).
• SEPTOGARD® suigtablette bevat bestanddele wat nie geskik is vir gebruik in kinders onder die ouderdom van 4 jaar nie.
• Indien u enige chroniese medikasie gebruik, moet nie hierdie produk gebruik sonder om u mediese praktisyn te raadpleeg nie.

Waarskuwings en voorsorgmaatreëls
Noukeurige sorg behoort geneem te word met SEPTOGARD® suigtablette.
• Raadpleeg 'n gesondheidsorgverskaffer of dokter voor gebruik indien u enige van die volgende het/neem:
• 'n bloedversteuring of bloedverdunners (antikoagulant- of antiplaatjie-medikasie) (sien **Ander medikasie saam met SEPTOGARD® suigtablette**); of
• ('n geskiedenis) van galstene, galbuis obstruksie, maagulkusse of oormaat maagsuur, aangesien SEPTOGARD® suigtablette borrie-risoom bevat.
• Indien u dokter u ingelig het dat u 'n onverdraagsaamheid teenoor sekere suikers het, moet u eers u dokter raadpleeg voordat u SEPTOGARD® suigtablette gebruik.
• SEPTOGARD® suigtablette bevat glukose en sukrose wat 'n effek op die beheer van u bloedsuikervlakke kan hê indien u diabetes mellitus het.
• Staak gebruik en raadpleeg 'n gesondheidsorgverskaffer of dokter indien u seerkeel simptome vererger of voortduur vir meer as 2 dae.
• Staak gebruik en raadpleeg 'n gesondheidsorgverskaffer of dokter indien u hoes vererger of aanhou vir meer as 7 dae of gepaard is met koors.
• Staak gebruik teen minste 2 weke voor enige chirurgiese prosedures.
• Moet nie die aanbevole daaglikse dosis oorskry nie.
• Moet nie na die vervaldatum gebruik nie.
• Produkte wat kruiebestanddele bevat kan interaksies met sommige medikasie hê.
Porfirie: Veiligheid is nog nie vasgestel nie.
U moet u dokter raadpleeg indien u onseker is van enige van die bogenoemde.

Ander medikasie en SEPTOGARD® suigtablette
Lig altyd u gesondheidsorgverskaffer in wanneer u medisyne op 'n gereelde basis neem, insluitend komplementêre- of tradisionele medisyne.
Raadpleeg u dokter, apteker of gesondheidsorgverskaffer voor gebruik indien u die volgende medisyne neem:
• medikasie vir hoë bloeddruk, aangesien SEPTOGARD® suigtablette bestanddele bevat wat die doeltreffendheid van die middels kan beïnvloed;
• bloedverdunners (antikoagulant- of antiplaatjie-medikasie) bv. warfarin, aangesien SEPTOGARD® suigtablette bestanddele bevat wat die effektiwiteit van die middels kan beïnvloed;
• hartglikosiede (hartmedikasie) bv. digoksien, aangesien dropwortel die risiko van kardiotoksisiteit van verhoog;
• kortikosteroïede, aangesien dropwortel die effekte en newe effekte van kortikosteroïede kan verhoog;
• sulfasalasien (vir artritis), aangesien borrie-risoom die effekte en newe effekte van sulfasalasien kan verhoog; en
• thiazied-diuretikums (waterpille), stimulerende lakseermiddels of medikasie wat elektrolietbalans kan beïnvloed, aangesien dropwortel kalium verlies kan veroorsaak.

Gebruik van SEPTOGARD® suigtablette saam met kos, drinkgoed en alkohol
Daar is geen bekende interaksie tussen die inhoud van SEPTOGARD® suigtablette en kos, drinkgoed of alkohol nie.

Swangerskap, borsvoeding en vrugbaarheid
Moet nie SEPTOGARD® suigtablette gebruik terwyl u swanger is of borsvoed nie, aangesien dit dropwortel en beleriese myrobalan-vrug bevat, wat teenaangedui is. Indien u swanger is of borsvoed, dink u is dalk swanger of beplan om 'n baba te hê, raadpleeg asseblief u dokter, apteker of ander gesondheidsorgverskaffer vir advies voordat u hierdie produk gebruik.

Bestuur en die gebruik van masjiene
SEPTOGARD® suigtablette behoort nie u vermoë om te bestuur of masjinerie te gebruik, te beïnvloed nie. Nietemin, is die nie altyd moontlik om te voorspel tot watter mate SEPTOGARD® suigtablette kan inmeng met u daaglikse aktiwiteite nie. U moet verseker dat u nie by bogenoemde aktiwiteite betrokke raak voordat u bewus is van die mate waarin SEPTOGARD® suigtablette u affekteer nie.

SEPTOGARD® suigtablette bevat glukose en sukrose
Indien u dokter u ingelig het dat u 'n onverdraagsaamheid teenoor sekere suikers het, moet u eers u dokter raadpleeg voordat u SEPTOGARD® suigtablette gebruik.
3. Hoe om SEPTOGARD® suigtablette te gebruik
Moet nie u medisyne met enige ander persoon deel nie. Gebruik SEPTOGARD® suigtablette altyd presies soos voorgeskryf in hierdie brosjure of soos u dokter, apteker of verpleegster u aanbeveel het. Maak asseblief seker by u dokter, apteker of verpleegster wat die aanbevole dosis is indien u onseker is.

Die gewone dosis vir volwassenes en kinders 4 jaar en ouer is: Los een (1) suigtablet stadig, elke drie tot vier ure, in die mond op. Moet nie meer as ses (6) suigtablette per 24-uur gebruik nie. Vir beste resultate, moet nie enigiets eet of drink binne 10 minute nadat suigtablet geneem is nie.

Indien u meer SEPTOGARD® suigtablette gebruik het as wat u moes
In die geval van 'n oordosis, raadpleeg u dokter of apteker. Indien nie een van hulle beskikbaar is nie, kontak die naaste hospitaal of gifbeheersentrum.

Indien u vergeet om SEPTOGARD® suigtablette te gebruik
Gebruik die dosis wat oorgeslaan is, sodra u daarvan onthou. Indien dit egter amper tyd vir die volgende dosis is, slaan die vergete dosis oor en gebruik SEPTOGARD® suigtablette op die volgende geskeduleerde tyd. Moet nie 'n dubbele dosis gebruik om te vergoed vir individuele dosisse wat u oorgeslaan het nie.

Wanneer u ophou om SEPTOGARD® suigtablette te gebruik
Indien u die dosis gebruik soos aangedui, sal daar waarskynlik geen ontrekkingseffek wees wanneer u die gebruik van hierdie medikasie staak nie.

4. Moontlike newe-effekte
SEPTOGARD® suigtablette kan newe-effekte hê. Nie alle newe-effekte wat aangemeld is vir SEPTOGARD® suigtablette is in hierdie brosjure ingesluit nie. Indien u algemene gesondheid vererger, of indien u ongewenste gevolge ervaar tydens die gebruik van SEPTOGARD® suigtablette, raadpleeg u gesondheidsorgverskaffer vir advies.

Indien een van die volgende of ander newe-effekte voorkom, stop die gebruik van hierdie produk en raadpleeg u dokter of apteker of gaan na die naaste ongevalle-afdeling by u naaste hospitaal:
Allergiese reaksies soos: veluitslag, knoppe (velbultjies), jeukerigheid, swelling van die gesig, lippe, mond of keel, wat probleme met sluk of asemhaling kan veroorsaak. Hierdie is almal baie ernstige newe-effekte. Indien u dit ervaar, het u dalk 'n ernstige reaksie op SEPTOGARD® suigtablette gehad. U mag dalk dringende mediese aandag of hospitalisasie benodig.

Vertel u dokter of apteker indien u enige van die volgende waarneem:
Frekwensie nie-bekend:
• Hoofpyn.
• Buikpyn of -ongemak, anale brandende sensasie, winde opbreek, opgeblasenheid, brandende nasmaak, hardlywigheid, diarree, droëmond, slegte spysvertering, sooibrand, naarheid en braking.

Lig asseblief u dokter of apteker in indien u enige newe-effekte opmerk wat nie in hierdie brosjure genoem word nie.

Rapportering van newe-effekte
Praat met u dokter, apteker, of verpleegster indien u newe-effekte ervaar. U kan ook newe-effekte by SAHPRA aanmeld via die Med Safety TOEP (Medsafety X SAHPRA) en eRapportering platvorm (who-umc.org), beskikbaar op SAHPRA se webwerf. Deur newe effekte aan te meld kan u help om meer inligting rakende die veiligheid van SEPTOGARD® suigtablette te verskaf.

5. Hoe om SEPTOGARD® suigtablette te bewaar
• Bewaar alle medisyne buite die bereik van kinders.
• Bewaar in 'n droë plek by of benede 25 °C.
• Beskerm teen lig en vog.
• Hou in oorspronklike verpakking totdat gebruik word.
• Moet nie in die yskas verkoel nie.
• Moet nie in die badkamer stoor nie.
• Moet nie ná die vervaldatum op die verpakking gebruik nie.
• Neem alle ongebruikte medisyne terug na u apteker.
• Moet nie ongebruikte medisyne in dreine of rioolstelsels (bv. toilette) weggooi nie.

6. Inhoud van die verpakking en ander inligting
Wat SEPTOGARD® suigtablette bevat

Elke suigtablet bevat:	Per suigtablet	Per maksimum daaglikse dosis (6 suigtablette)
Mentha piperata (peperment/Sanskrit – Pudina Ke Phool) [kristalle]	7 mg	42 mg
Uittreksel van die volgende:		
Glycyrrhiza glabra L. (drop/Sanskrit - Yashtimadhu) [wortel]	20 mg	120 mg
Terminalia belerica Roxb. (beleriese myrobalan/Sanskrit - Baheda) [vrug]	5,5 mg	33 mg
Zingiber officinale Rosc. (gemmer/Sanskrit – Sonth) [risoom]	4,5 mg	27 mg
Curcuma longa L. (borrie/Sanskrit – Haldi) [risoom]	2 mg	12 mg

Die ander bestanddele en hulpstowwe is: sitroensuur-monohidraat.
Heuning-Suurlemoen geur: heuning-suurlemoen geursel, kinolien-geel kleurstof.
Kruisement geur: propileenglikol, glansblou kleurstof, kinolien-geel kleurstof.
Lemoen geur: lemoen geursel, sonstraal-geel kleurstof.

SEPTOGARD® suigtablette bevat suiker: sukrose 1375 mg/suigtablet, vloeibare glukose 1375 mg/suigtablet.
SEPTOGARD® suigtablette bevat preserveermiddels: kaliumsorbaat 0,2 % m/m, sorbiensuur 0,02 % m/m.

Hoe SEPTOGARD® suigtablette lyk en die inhoud van die verpakking
SEPTOGARD® suigtablette is goudgeel, groen of oranje gekleurde, plat, afgeskuinste, ronde suigtablette met 'n metanoliese soet smaak, kenmerkend van heuning suurlemoen, kruisement of lemoen geur. SEPTOGARD® suigtablette is beskikbaar as 12 suigtablette (wat 2 blisterverpakkings van 6 suigtablette elk bevat) of 24 suigtablette (wat 4 blisterverpakkings van 6 suigtablette elk bevat) in 'n gedrukte eenheidskarton, wat 'n pasiëntinligtingsbrosjure insluit.

Houer van Sertifikaat van Registrasie
Nativa (Edms) Bpk
Cradocklaan 260, Lyttelton, Centurion, 0157, Gauteng, Suid-Afrika
Tel: +27 (0) 12 664 7110
Verbruikershulplyn: 0860 628 482
E-pos: health@nativa.co.za
Webwerf: www.nativa.co.za

Hierdie brosjure was laas hersien in
April 2025

Datum van registrasie
Moet toegeken word

Registrasienommer
Moet toegeken word

Professionele Inligting
Die SEPTOGARD® suigtablette Professionele Inligting kan verkry word op die Nativa webwerf (www.nativa.co.za).

PPAT155/02

PEI0465 0425 E

FRONT

BACK

INSERT SIZE : 195 x 275 mm
FOLDED SIZE : 49 x 69 mm
Folding Pattern : 2 Horizontal & 2 Vertical
PRINT COLOUR : BLACK

EXPORTS - SOUTH AFRICA
AT: LEAMAK

Reason for change:
Common insert for 3 flavours

Digitally signed by Annemie van Heerden
Date: 2025.06.02 16:37:52 +02'00'

MASTER

Digitally signed by Gail Young
Date: 2025.06.03 19:28:34 +02'00'